than five minutes. When dried, the sediment was homogenized and placed into wide mouth sample containers.

All of M&E's field sampling was conducted between November 12, 1997 and November 20, 1997. The weather during the sampling event was generally clear and cold with temperatures falling from the start of the event from the mid 40's°F to about 30°F at the end. Snow and sleet fell on November 14th, but sampling continued as there was no wind. Sample dewatering on that day took place under tarps to exclude falling precipitation from the samples.

### 2.2 Analytical Program

The analytical program used to analyze the sediment samples collected by M&E is described in the Final QAPP (M&E, 1997b). All analyses were conducted under the EPA DAS program following EPA Region I approved DAS analytical methodologies. The EPA-approved DAS analytical specifications are provided in Appendix B of the QAPP (M&E, 1997b). No changes to the methods were proposed from the results of the pre-test sampling. Analysis for VOCs, PAHs, pesticides, PCBs, and grain size were conducted by Southwest Research Institute in San Antonio, Texas. Nine of the sediment samples were reanalyzed for PAHs using Gas Chromatography/Mass Spectroscopy/Selected Ion Monitoring (GC/MS/SIM) due to low concentrations of PAHs, and in order to meet the required reporting limits for the ecological risk assessment (M&E, 1997a and 1997c). Analysis for low concentration metals, total combustible organics (TCO), moisture content, and pH were conducted by RECRA LabNet Philadelphia, in Lionville, Pennsylvania. The AVS/SEM analyses were conducted by American Environmental Network in Monroe, Connecticut. Table 2-3 summarizes laboratory information about the analyses that were conducted for the samples collected. No significant deviations occurred from the DAS analytical methods.

## 3.0 SEDIMENT SAMPLE RESULTS

Validated data for all sediment and field quality control (QC) samples collected during M&E's field investigation are presented in Appendix B. The data are presented by analytical fraction and method in sequential sample station/location order:

- VOCs
- PAHs
- PAHs GC/MS/SIM
- Pesticides
- PCBs
- Metals (total extractable)
- AVS/SEM
- TCO
- Grain size -- sieve
- Grain size -- hydrometer

Table 2-3
Sample Location and Laboratory Reference Table

Sample Local	T		QC	Case	DAS	SDG	SÞG	SDG
Sample	Date Collected	Matrix	Sample	Number	Number	Lab - SWRI	Lab - RECRA	Lab - IEA
Location		Sediment	Cumpic	0049M	DAM322	DAM320-OA	22.1.1.	DAM320-OB
SED-01-06	12-Nov-97			0049M	DAM323	DAM320-OA	DAM320-IA	DAM320-OB
SED-01-07	12-Nov-97			0049M	DAM326	DAM320-OA	DAM320-IA	DAM320-OB
SED-02-01	13-Nov-97			0049M	DAM327	DAM320-OA	DAM320-IA	DAM320-OB
SED-02-02		Sediment		0049M	DAM325	DAM320-OA	DAM320-IA	DAM320-OB
SED-03-02	13-Nov-97   14-Nov-97			0049M	DAM333	DAM320-OA	DAM320-IA	DAM320-OB
SED-04-02	14-Nov-97			0049M	DAM334	DAM320-OA	DAM320-IA	DAM320-OB
SED-04-03	14-Nov-97 13-Nov-97		Dunlicate	0049M	DAM328	DAM320-OA	DAM320-IA	DAM320-OB
SED-05-03	13-Nov-97 13-Nov-97			0049M	DAM331	DAM320-OA	DAM320-IA	DAM320-OB
SED-05-05		1	Dupiteate	0049M	DAM345	DAM340-OA	DAM345-IA	DAM345-OA
SED-06-03	18-Nov-97	1		0049M	DAM359	DAM340-OA	DAM345-IA	DAM345-OA
SED-07-02	20-Nov-97	Sediment	MC/MSD	0049M	DAM360	DAM340-OA	DAM345-IA	DAM345-OA
SED-07-05	20-Nov-97		MISHMOD	0049M	DAM353	DAM340-OA	DAM345-IA	DAM345-OA
SED-07-10	19-Nov-97		Dumlioata	0049IM	DAM351	DAM340-OA	DAM345-IA	DAM345-OA
SED-10-01	19-Nov-97	1	Duplicate Duplicate	0049M	DAM355	DAM340-OA	DAM345-IA	DAM345-OA
SED-10-04		Sediment	· "	0049M	DAM352	DAM340-OA	DAM345-IA	DAM345-OA
SED-10-02		Sediment		0049M	DAM336	DAM320-OA	DAM320-IA	DAM320-OB
SED-11-01		Sediment		0049M	DAM357	DAM340-OA	DAM345-IA	DAM345-OA
SED-12-01		Sediment		0049M	DAM358	DAM340-OA	DAM345-IA	DAM345-OA
SED-12-03		Sediment		0049M	DAM341	DAM340-OA	DAM320-IA	DAM320-OB
SED-13-01	17-Nov-97	1	l	0049M	DAM342	DAM340-OA		DAM320-OB
SED-13-03	17-Nov-9	1	1	0049M	DAM340	I	l'	DAM320-OB
SED-15-01	17-Nov-9	ı		0049M		1		DAM345-OA
SED-18-02	18-Nov-9		h .	0049M	1	1		DAM345-OA
SED-18-03		7 Sedimen		0049M	l l			DAM345-OA
SED-19-01	19-Nov-9	7 Sedimen	LACATE	1			1	DAM320-OB
SED-20-01	1		t MS/MSE				1	DAM320-OB
SED-21-01		7 Sedimen		1	1		·	
SED-21-05		7 Sedimen		l I	. h			DAM320-OB
SED-22-02		7 Sedimer		0049M		l l		DAM320-OB
SED-24-03		7 Sedimer		0049M		1	- 1	
SED-25-02	18-Nov-9	7 Sedimer	nt	0049M			1	
SED-12-TE	3   12-Nov-9	7 Methano	ol Trip Blan	k 0049M		1		Ì
SED-13-TE	3   13-Nov-9	97   Methan	ol Trip Blan	ık 0049M	1			
SED-14-TI	3 14-Nov-9	97   Methan	ol Trip Blar	ık 0049N				
SED-17-TI	B   17-Nov-9	97 Methan	ol Trip Blan	nk 0049N	1	1		
SED-18-TI	B 18-Nov-9	97 Methan	ol Trip Blai	nk 0049N	1	1		
SED-19-T	B   19-Nov-	97 Methan	ol Trip Bla	nk 0049N	L	1		
SED-20-T	B 20-Nov-	97 Methan	ol Trip Bla	nk 0049N				DAM320-OB
SED-13-P	E 13-Nov-	97 Sedime	nt PE Samp	ole   0049N	1	1	· · }	
SED-18-P	L	97 Sedime	nt PE Samp	ole 0049N	I	1		l l
SED-17-E		97 Aqueo	us Equip. E	sik  00491			1	
SED-20-E	i i		us Equip. E		и DAM36	2 DAM340-O	A DAM343-17	A DEMINISTRA

# 3.1 Analytical Data Validation

Quality control (QC) samples were taken and submitted for laboratory analysis to monitor precision, accuracy, and potential contamination throughout the sampling episode. These samples included trip blanks, equipment blanks, field duplicates, and matrix spike/matrix spike duplicate samples.

All sediment analytical data generated from field investigation were validated to EPA Region I's Tier III level in accordance with Region I EPA New England Data Validation Guidelines (U.S. EPA, 1996), which were modified to meet the QC criteria set forth in the DAS analytical specifications. Typical actions implemented during validation of analytical data are described in Appendix D and are summarized below.

Data found to be suspect during the validation process were qualified, and a description of the discrepancy was noted in the validation summary. Positive and non-detect results that were shown to have serious QC problems were rejected and flagged with an "R." Positive results that were shown to exhibit poor precision or accuracy were qualified as approximated and flagged with a "J." Nondetect results that were shown to exhibit poor precision were flagged with a "UJ." Nonqualified positive results were found to meet all validation criteria. Nondetect results that were found to meet the validation criteria were shown as the quantitative limit or detection limit followed by a "U" qualifier. Rejected values were considered as non-usable results for subsequent evaluations. Approximated and nonqualified results were used in further evaluations, but the qualified data were first reviewed to establish their usability.

The data tables presented in Appendix B show the results of the validation. As was described in Section 3.1, the % solids in each sediment sample were increased (the moisture content decreased) using field techniques and laboratory procedures specifically designed for this project. All results presented in the Appendix B data tables are in dry weight units; however, the % moisture for each sample is listed at the bottom of the tables. This value represents the % moisture after field manipulation. Only sediment samples collected for VOC analysis were analyzed at the "true" or unmanipulated moisture content found in the field.

## 3.2 Sediment Data Results

Table 3-1 summarizes by station/location the validated data for analytes that were detected in at least one sediment sample collected by M&E.

## 4.0 DATA EVALUATION

This section discusses the comparison between the 1995 and 1997 sediment data as well as data trends for the major chemical groups detected across the site.

TABLE 3-1. ANALYTES DETECTED IN 1997 SEDIMENT SAMPLES (1,2)

M&E SAMPLE ID: DAS SAMPLE NUMBER:	SED-01-06 DAM322	SED-01-07 DAM323	SED-02-01 DAM326	SED-02-02 DAM327	SED-03-02 DAM325	SED-04-02 DAM333	SED-04-03 DAM334 11/14/97
DATE SAMPLED:	11/12/97	11/12/97	11/13/97	11/13/97	11/13/97	11/14/97	11/14/9/
REMARKS:							
OCs (ug/Kg)				21 111	9 J	24 UJ	32 UJ
Benzene	26 UJ	30 UJ	10 UJ	21 UJ 53 UJ	44 UJ	60 UJ	224 UJ
Chloroform	66 UJ	76 UJ	26 UJ		28 J	60 UJ	74 UJ
cis-1,2-Dichloroethene	62 J	58 J	26 UJ	53 UJ	18 UJ	24 UJ	29 UJ
Ethylbenzene	26 UJ	30 UJ	10 UJ	21 UJ	44 UJ	60 UJ	74 UJ
m/p-Xylene	66 UJ	76 UJ	26 UJ	53 UJ	132 UJ	180 U	221 U
Naphthalene	197 UJ	228 UJ	77 UJ	159 UJ		180 U	221 U
Tetrachloroethene	197 UJ	228 ÚJ	77 UJ	159 UJ	132 UJ 132 UJ	. 180 U	221 U
rans-1,2-Dichloroethene	197 UJ	228 UJ	77 UJ	159 UJ		60 UJ	74 UJ
Trichloroethene	37 J	76 UJ	26 UJ	53 UJ	44 UJ	180 U	221 U
Vinyl chloride	1 <b>9</b> 7 UJ	228 UJ	77 UJ	159 UJ	132 UJ	180 0	221 0
PAHs (ug/Kg)				20. 1	16 J	41 J	70
2-Methylnaphthalene	20 J	20 J	10 J	20 J 30 J	45 J	110 J	170 J
Acenaphthene	33 J	32 J	R	57 J	39 J	170	160
Acenaphthylene	33 J	37 J	19 J		120	420	640
Anthracene	92	100	62 J	140 1500	1300	3100	3200
Benzo(a)anthracene	1000	1100	590	1400	1000	4100	3500
Benzo(a) pyrene	930	920	620		1400	5100	6000
Benzo(b)fluoranthene	1400	1600	720	1700	570	1900	2100
Benzo(g,h,i)perylene	480	510	330	640	640	2100	3800
Benzo(k)fluoranthene	730	690	340	710	1300	3700	3800
Chrysene	880	980	660	1400		720	890
Dibenz(a,h)anthracene	120	110	60 J	150	110	8300	8500
Fluoranthene	2100	2200	1200	3000	2600	290	490
Fluorene	89	89	44 J	97	150	2600	2500
Indeno(1,2,3-cd)pyrene	580	640	420	860	620	2000 56 J	96 J
Naphthalen <del>e</del>	24 J	23 J	13 J	24 J	18 J	3200	3400
Phenanthrene	840	900	390	970	1000		6000
Рутепе	1900	1900	1100	2500	2600	6700	0000
PAH-SIMS (ug/Kg)							
Naphthalene			13 UJ				
2-Methylnaphthalene			12 J				
Acenaphthylene			22 J 11 J				
Acenaphthene			11 J 42 J				
Fluorene			69 J				
Dibenz(a,h)anthracene			<b>4</b> 2 <b>4</b>				

TABLE 3-1. ANALYT ETECTED IN 1997 SEDIMENT SAMPLES (1,2)

M&E SAMPLE ID:	SED-01-06	SED-01-07	SED-02-01	SED-02-02	SED-03-02	SED-04-02 DAM333	SED-04-03 DAM334
DAS SAMPLE NUMBER:	DAM322	DAM323	DAM326	DAM327	DAM325		11/14/97
DATE SAMPLED:	11/12/97	11/12/97	11/13/97	11/13/97	11/13/97	11/14/97 	11/17/2/
REMARKS:							
esticides (ug/Kg)			20. 1	25 J	94	160	37 J
,4'-DDD	16 J	18 J	20 J	30	29	91	20 J
,4'-DDE	24	28	26 J	2.4 J	5.8 J	21 J	4.4 J
,4'-DDT	2.5 J	3.2 J	1.8 J	0.16 U	1.6 U	1.9 U	0.30 J
ldrin	0.16 U	0.21 J	0.16 UJ		1.6 U	1.9 U	0.16 U
lpha-BHC	0.16 U	0.17 J	0.16 UJ	0.16 U	1.0 U	21	29 J
Ipha-Chlordane	9.3 J	10 J	11 J	16	2.3 J	1.9 U	1.3 J
elta-BHC	0.19 J	0.16 U	1.3 J	0.16 U		3.9 U	0.73 J
Dieldrin	0.50 J	0.61 J	0.45 J	0.47	3.2 U	1.9 U	0.16 U
Indosulfan I	0.16 U	0.16 U	0.16 UJ	0.17 U	1.6 U	3.9 U	0.49
Indosulfan II	0.41	0.66	0.21 J	0.28	3.2 U	3.9 U	0.33 J
Endosulfan Sulfate	0.67 J	0.78 J	0.47 J	0.33 U	3.2 U		
Endrin	1.2 J	1.8 J	1.3 J	1.9	3.2 U	5.0 J	3.2 1
endrin Aldehyde	3.4 J	3.2 J	1.6 J	1.9 J	3.2 U	11 <b>J</b>	4.9 1
gamma-BHC	0.16 U	0.16 U	0.16 UJ	0.16 U	1.6 U	1.9 U	0.16
ramma-Chlordane	6.4 J	6.8 J	8.0 J	11 J	9.3 J	16 J	22
Heptachlor Epoxide	0.16 U	0.16 U	0.16 UJ	0.16 U	1.6 U	1.9 U	0.16
PCBs (ug/Kg)					,	200 1	44
Aroclor-1248	46 J	70 J	160 J	180 J	160 J	200 J	160
Aroclor-1260	47	69 J	140	170	110	94	(00)
Total Metals (mg/Kg)					6720 J	17700 J	11500
Aluminum	12200 J	13200 J	6570 J	11200 J	1.7 J	11.1 J	1.8
Antimony	1.7 J	1.8 J	1.9 J	2.1 J	86.5	1570	98.6
Arsenic	78.1	111	56.4	105	84.6	77.3	175
Barium	85.8	95.1	50.6	81.2	0.59 J	1.7 J	0.86
Beryllium	0.87 J	0.98 J	0.47 J	0.91 J	5.2	19.1	7.4
Cadmium	6.8	8.5	3.8	16.1 4600	4410	5900	6310
Calcium	5420	7510	2610	4000 190 J	200 J	3000 J	240
Chromium	148 J	192 J	103 J	33.0 J	21.4 J	42.7 J	21.6
Cobalt	27.9 J	33.1 J	17.0 J	381	171	1080	323
Copper	305	436	128	26000 J	21900 J	86200 J	31600
Iron	31800 J	32500 J	16900 J	2000 J 449 J	345 J	425 J	487
Lead	413 J	477 J	225 J	2510	1790	1400	3470
Magnesium	3480	3430	2150	390	503	803	1010
Manganese	479	595	206	1.1	0.71	13.6	1.7
Mercury	1.0	1.1	0.37 21.5 J	31.0 J	26.7 J	41.2 J	33.6
Nickel	40.4 J	41.3 J	21.5 J 640 J	784 J	604 J	476 J	1120
Potassium	[190 J	1170 J	040 J	707 3			

TABLE 3-1. ANALYTES DETECTED IN 1997 SEDIMENT SAMPLES (1,2)

M&E SAMPLE ID:  DAS SAMPLE NUMBER:  DATE SAMPLED:  REMARKS:	SED-01-06	SED-01-07	SED-02-01	SED-02-02	SED-03-02	SED-04-02	SED-04-03
	DAM322	DAM323	DAM326	DAM327	DAM325	DAM333	DAM334
	11/12/97	11/12/97	11/13/97	11/13/97	11/13/97	11/14/97	11/14/97
elenium	5.1 J	1.4 J	1.6 J	2.0 J	4.2 J	1.8 J	2.0 J
ilver	0.75 J	0.66 J	0.40 J	0.91 J	0.69 J	0.59 J	0.91 J
Sodium	583	611	197	382	321	330	347
Fhallium	0.63 UJ	0.43 UJ	0.53 UJ	0.48 UJ	0.43 UJ	0.44 UJ	0.43 UJ
Vanadium	56.4	55.3	32.2	46.1	49.2	63.8	55.7
Linc	2080 J	2370 J	992 J	5170 J	1420 J	4330 J	1590 J
AVS-SEM (umoles/g) Arsenic Cadmium Copper Lead Nickel Sulfide Zinc SEM/AVS Ratio	0.06 J 0.08 UJ 0.05 UJ 1.87 J 0.86 J 72.0 26.5 J	0.11 J 0.09 J 0.08 J 2.13 J 0.23 J 79.9 29.3 J 0.40	0.05 U 0.05 U 0.05 UJ 0.83 J 0.14 J 34.7 11.2 J 0.35	0.05 J 0.15 J 0.05 UJ 1.88 J 0.49 J 129 36.7 J 0.30	0.05 U 0.08 UJ 0.05 UJ 1.69 J 0.98 J 88.0 25.0 J 0.32	0.09 J 0.08 UJ 0.05 UJ 1.96 J 0.24 J 23.5 21.0 J	0.08 J 0.06 U 0.05 U 1.60 J 0.34 J 18.7 15.1 J 0.91
TCO (mg/Kg) Total Combustible Organics	230000 J	152000 J	125000 J	160000 J	195000 J	193000 J	255000 J

(1) Validated data for all sediment and field QC samples collected during M&E's field investigation are reported in Appendix B.

(2) Data presented for an analyte if the analyte was detected in at least one sediment sample

- J Quantitation is approximate due to limitations identified in the quality control review
- U Value reported is the sample-specific detection limit
- R Value is rejected
- UJ Sample-specific detection limit is approximate due to limitations identified in the quality control review

TABLE 3-1. ANALYTE DETECTED IN 1997 SEDIMENT SAMPLES (1,2)

M&E SAMPLE ID: DAS SAMPLE NUMBER: DATE SAMPLED:	SED-05-03 DAM328 11/13/97	SED-05-05 DAM331 11/13/97	SED-06-03 DAM345 11/18/97	SED-07-02 DAM359 11/20/97	SED-07-05 DAM360 11/20/97	SED-07-10 DAM353 11/19/97 	SED-10-01 DAM351 11/19/97 FD of SED-10-04
REMARKS:  OCs (ug/Kg) enzene Chloroform is-1,2-Dichloroethene Cthylbenzene n/p-Xylene Naphthalene Tetrachloroethene Trichloroethene Trichloroethene Vinyl chloride PAHs (ug/Kg)	6 J 9 UJ 7 J 5 J 10 J 17 J 28 U 28 U 6 J 28 U	FD of SED-05-03	17 J 32 UJ 38 J 9 J 25 J 70 J 96 U 96 U 52 J 96 U	10 UJ 25 UJ 20 J 5 J 25 UJ 74 UJ 74 UJ 16 J 74 UJ	11 UJ 26 UJ 18 J 11 UJ 26 UJ 79 UJ 79 UJ 24 J 79 UJ	13 UJ 34 UJ 24 J 13 UJ 34 UJ 101 UJ 101 UJ 35 J 101 UJ	44 UJ 110 UJ 110 UJ 44 UJ 110 UJ 330 U 330 U 330 U 330 U 37 J 27 J
2-Methylnaphthalene Acenaphthene Acenaphthylene Anthracene Benzo(a)anthracene Benzo(a)pyrene Benzo(b)fluoranthene Benzo(g,h,i)perylene Benzo(k)fluoranthene Chrysene Dibenz(a,h)anthracene Fluoranthene Fluoranthene Fluorene Indeno(1,2,3-cd)pyrene Naphthalene Phenanthrene Pyrene PAH-SIMS (ug/Kg)	41 J 84 J 30 J 200 830 660 770 280 67 UJ 890 95 1900 160 J 340 34 J 1300 J 1600	70 120 J 42 J 290 1200 810 700 330 860 J 980 130 2600 290 J 370 35 J 2200 J 2600	130 J 120 550 4200 J 5100 J 6100 J 1900 J 1300 J 6900 J 720 J 9900 400 2400 J 53 J 5200 7500	220 J 140 740 6000 6000 8800 J 3300 4100 J 6600 1200 J 14000 640 4200 61 J 8500 9500	170 J 130 880 5200 5700 7800 J 3400 1800 J 6000 1400 J 14000 500 4200 57 J 4800 9800	230 J 140 890 9600 10000 16000 J 5300 2300 J 10000 2000 23000 550 6900 72 J 12000 15000	16 J 93 500 540 830 : 330 470 520 110 1600 110 360 86 600

Naphthalene 2-Methylnaphthalene Acenaphthylene

Acenaphthene

Fluorene

Dibenz(a,h)anthracene

TABLE 3-1. ANALYTES DETECTED IN 1997 SEDIMENT SAMPLES (1,2)

M&E SAMPLE ID: DAS SAMPLE NUMBER: DATE SAMPLED: REMARKS:	SED-05-03 DAM328 11/13/97 FD of SED-05-05	SED-05-05 DAM331 11/13/97 FD of SED-05-03	SED-06-03 DAM345 11/18/97	SED-07-02 DAM359 11/20/97	SED-07-05 DAM360 11/20/97	SED-07-10 DAM353 11/19/97	SED-10-01 DAM351 11/19/97 FD of SED-10-04
Pesticides (ug/Kg)				· · · · · · · · · · · · · · · · ·		36	98 J
-esuciues tug/ise/	180	180	79	47	39 J	33	90 J
,4'-DDE	37	34	71	19 J	41	14 J	3.4 U
	9.7 J	8.8 J	15 J	21 J	18 J		11 J
1,4'-DDT	1.7 U	1.7 U	1.6 U	1.6 U	1.6 U	1.6 U	1.7 U
Aldrin	1.7 U	1.7 U	1.6 U	1.6 U	1.6 U	1.6 U	
llpha-BHC	9.1 J	17 J	80	49 J	64	80	18 J
llpha-Chlordane	• •	4.3 J	25 J	1.6 U	24 J	1.6 U	1.7 U
lelta-BHC	3.0 J	3.3 U	5.0 J	4.1 J	3.2 U	3.2 U	13
Dieldrin	4.2 J		1.6 U	1.6 U	1.6 U	1.6 U	1.7 U
Endosulfan I	1.7 U	1.7 U	3.2 U	3.3 U	3.2 U	3.2 U	3.4 t
Endosulfan II	3.3 U	3.3 U		3.3 U	3.2 U	3.2 U	3.4 t
Endosulfan Sulfate	3.3 U	3.3 U	3.2 U	8.6 J	17 J	16 J	4.7 ]
Endrin	3.3 U	3.3 U	17 J	27 J	3.2 U	3.2 U	6.4 J
Endrin Aldehyde	3.3 U	3.3 U	3.2 U		1.6 U	1.6 U	1.7
gamma-BHC	1.7 U	1.7 U	1.6 U	1.6 U	38 J	45 J	16 )
gamma-Chlordane	7.3 J	16 J	48 J	41		1.6 U	1.7
Heptachlor Epoxide	1.7 U	1.7 U	1.6 U	1.6 U	1.6 U	1.0 0	,,,,
PCBs (ug/Kg)		_	120 1	37 J	29 J	67 J	300 J
Aroclor-1248	120 J	120 J	120 J		74	50	190
Aroclor-1260	77	72	82	95	74		
Total Metals (mg/Kg)		4000 T	14300	13200	14700	12500	30100
Aluminum	4860 J	4300 J	2.5 J	2.6 J	2.8 J	2.4 J	17.7
Antimony	2.4 J	0.68 UJ		111	98.3	87.4	715
Arsenic	9.8	9.9	135 130	110	118	89.8	99.2
Barium	40.8	37.8	1.0	0.95	1.0	0.86	2.0
Beryllium	0.32 J	0.27 J	7.7	6.3	5.8	4.5	17.5
Cadmium	0.76	0.48	5510	5310	6630	5840	9570
Calcium	1590	1090	403	291	277	238	3290
Chromium	90.0 J	46.9 ]	403 20.9	19.2	21.5	18.2	34.5
Cobalt	5.6 J	4.4 J	20.9 389	306	304	261	2590
Copper	45.9	40.4	40600	33800	35400	29100	45500
Iron	9520 J	8600 J	40600 455	382	406	326	975
Lead	1090 J	220 J	433 4260	4180	5330	4420	2130
Magnesium	2100	1590		517	495	397	570
Manganese	103	91.9	1630	2.3	2.0	1.6	24.4
Mercury	0.23	0.22	2.6	35.2	37.5	31.5	41.4
Nickel	11.2 J	10.8 J	32.2	1310	1620	1320	597
Potassium	681 J	390 J	1380	1310	1020		

TABLE 3-1. ANALYTES DETECTED IN 1997 SEDIMENT SAMPLES (1,2)

TCO (mg/Kg) Total Combustible Organics	34200 J	84000 J	197000 J	194000 J	210000 J	218000 J	471000 J
∟ead Nickel Sulfide Zinc SEM/AVS Ratio	1.17 J 0.10 J 24.7 3.88 J 0.23	0.78 J 0.06 J 15.3 2.24 J 0.20	1.54 0.39 J 5.44 11.2 J 2.44	0.47 J 20.4 15.1 J 0.84	0.30 J 49.6 17.1 J 0.38	0.25 J 5.22 11.2 J 2.53	0.34 J 64.3 24.1 J 0.41
AVS-SEM (umoles/g) Arsenic Cadmium Copper	0.05 U 0.05 U 0.46 J	0.05 U 0.05 U 0.05 UJ	0.17 J 0.06 0.08	0.19 J 0.05 U 0.19 1.46	0.10 J 0.05 0.05 U 1.41	0.17 J 0.07 0.05 U 1.67	1.56 J 0.11 0.10 1.98 J
lenium lver odium hallium anadium inc	0.53 U 0.18 J 103 0.04 UJ 26.3 144 J	0.53 U 0.73 J 85.3 0.04 UJ 20.8 126 J	3.4 J 0.18 J 617 U 0.77 U 60.2 1630	2.2 J 0.86 J 416 U 0.81 U 56.7 1500	0.91 J 519 U 0.44 U 62.6 1450	0.78 J 483 U 0.76 U 54.6 1190	0.62 J 726 U 1.7 U 75.5 3340
M&E SAMPLE ID: DAS SAMPLE NUMBER: DATE SAMPLED: REMARKS:	DAM328 11/13/97 FD of SED-05-05	SED-05-05 DAM331 11/13/97 FD of SED-05-03	SED-06-03 DAM345 11/18/97	SED-07-02 DAM359 11/20/97	SED-07-05 DAM360 11/20/97 	SED-07-10 DAM353 11/19/97	SED-10-01 DAM351 11/19/97 FD of SED-10-04

(1) Validated data for all sediment and field QC samples collected during M&E's field investigation are reported in Appendix B.

(2) Data presented for an analyte if the analyte was detected in at least one sediment sample

- J Quantitation is approximate due to limitations identified in the quality control review
- U Value reported is the sample-specific detection limit
- R Value is rejected
- UJ Sample-specific detection limit is approximate due to limitations identified in the quality control review

TABLE 3-1. ANALYTES DETECTED IN 1997 SEDIMENT SAMPLES (1,2)

M&E SAMPLE ID:	SED-10-02	SED-10-04	SED-11-01	SED-12-01	SED-12-03	SED-13-01	SED-13-03
DAS SAMPLE NUMBER:	DAM352	DAM355	DAM336	DAM357	DAM358	DAM341	DAM342
DATE SAMPLED:	11/19/97	11/19/97	11/14/97	11/20/97	11/20/97	11/17/97	11/17/97
REMARKS:		FD of SED-10-01					
VOCs (ug/Kg)							
Benzene	21 J	42 UJ	23 UJ	41 UJ	9 J	8 J	7 J
Chloroform	93 UJ	104 UJ	154 UJ	103 UJ	41 UJ	35 UJ	32 UJ
cis-1,2-Dichloroethene	93 UJ	64 J	170 J	417 J	25 J	35 UJ	23 J
Ethylbenzene	37 UJ	42 UJ	23 UJ	41 UJ	16 UJ	9 J	13 UJ
m/p-Xylene	93 UJ	104 UJ	57 UJ	103 UJ	41 UJ	35 UJ	32 UJ
Naphthalene	278 U	312 U	171 U	309 U	124 U	104 UJ	97 UJ
Tetrachloroethene	278 U	312 U	171 U	<b>309</b> U	124 U	108 UJ	97 UJ
trans-1,2-Dichloroethene	278 U	312 U	387	309 U	124 U	104 UJ	97 UJ
Trichloroethene	60 J	104 UJ	103 UJ	53 J	41 UJ	35 UJ	37 J
Vinyl chloride	278 U	312 U	171 U	309 U	124 U	104 UJ	97 UJ
PAHs (ug/Kg)							
2-Methylnaphthalene	67 U	29 J	13 J	12 J	22 J	73	23 J
Acenaphthene	R	31 J	R	R	R	89 J	22 J
Acenaphthylene	8 J	24 J	9 J	16 J	14 J	31 J	26 J
Anthracene	49 J	100 J	54 J	69 J	79	300	99
Benzo(a)anthracene	210	660	410	560 J	420	1700	750
Benzo(a)pyrene	250	700	520	1000 J	440	1700	890
Benzo(b)fluoranthene	290 J	1100 J	780	1500 J	780 J	1900 J	1100 J
Benzo(g,h,i)perylene	200	400	300	640 J	290	820	540
Benzo(k)fluoranthene	220 J	430 J	720	720 J	440 J	850 J	570 J
Chrysene	380	780	670	850 J	610	1700	1100
Dibenz(a,h)anthracene	69	140	99	220 J	110	220	180
Fluoranthene	570	2300 J	1000	1200 J	750	5200	2300
Fluorene	59 J	120 J	46 J	47 J	59 J	220	67 J
Indeno(1,2,3-cd)pyrene	190	450	360	740 J	330	1000	610
Naphthalene	26 J	56 J	29 J	26 J	98 J	67 J	40 J
Phenanthrene	210	780 J	260	570 J	460	2700	960
Pyrene	470	2000 J	950	1000 J	650	4000	1700
PAH-SIMS (ug/Kg)							
Naphthalene	30 UJ		32 J	24 J	120 J		
2-Methylnaphthalene	10 U		16 J	13 J	25 J		
Acenaphthylene	13 J		9 J	19 J	23 J		
Acenaphthene	13		8 J	10 J	15 J		
Fluorene	72 J		57 J	45 J	78 J 150		1
Dibenz(a,h)anthracene	42 J		97 J	320 J	130		

TABLE 3-1. ANALYTES DETECTED IN 1997 SEDIMENT SAMPLES (1,2)

	SED-10-02	SED-10-04	SED-11-01	SED-12-01	SED-12-03	SED-13-01	SED-13-03
M&E SAMPLE ID:		DAM355	DAM336	DAM357	DAM358	DAM341	DAM342
DAS SAMPLE NUMBER:	DAM352	11/19/97	11/14/97	11/20/97	11/20/97	11/17/97	11/17/97
DATE SAMPLED:	11/19/97	FD of SED-10-01			••	****	
REMARKS:		TD 01 3LD-10-01					
esticides (ug/Kg)	4 #	260 J	45 J	17	1.9	18 J	310
,4'-DDD	17		15 J	4.0	0.90 J	17 J	130
,4'-DDE	5.1	230 J	1.9 J	1.3 J	1.1 J	4.0 J	12 J
,4'-DDT	0.35 U	5.3 J	, 0.16 UJ	0.17 U	0.16 U	0.16 UJ	1.6 U
Aldrin	0.17 U	25 J		0.17 U	0.16 U	0.16 UJ	1.6 t
lpha-BHC	0.17 U	1.7 U	0.16 UJ	2.4 J	1.1 J	17 J	93
dpha-Chlordane	1.5 J	41 J	8.1 J	0.17 U	0.16 U	0.16 UJ	15 J
lelta-BHC	0.17 U	1.7 U	0.16 UJ		0.32 U	1.4 J	11 /
Dieldrin	0.35 U	15 J	1.4 J	0.78 J	0.16 U	0.16 UJ	1.6
Endosulfan I	0.17 U	1.7 U	0.44 J	0.17 U		0.39 J	3.3
Endosulfan II	0.35 U	3.3 U	0.51 J	0.33 U	0.32 U	0.39 J 0.38 J	3.3 t
Endosulfan Sulfate	0.35 U	3.3 U	0.72 J	0.33 U	0.32 U		3.3
Endrin	0.61 J	6.9	0.69 J	1.4 J	0.77 J	9.6 J	7.5
Endrin Aldehyde	1.7 J	17 J	1.7 J	1.8 J	0.87 J	2.2 J	
zamma-BHC	0.17 U	1.7 U	0.16 UJ	0.23 J	0.34 J	0.16 UJ	1.6
gamma-Chlordane	0.25 J	25 J	5.6 J	1.7 J	0.73 J	16 J	650 .
Heptachlor Epoxide	0.54 J	2.3 Ј	0.16 UJ	0.46 J	0.27 J	0.40 J	1.6
•							
PCBs (ug/Kg)	47 J	820 J	180 J	28 J	15 J	25 J	170 .
Aroclor-1248		450 J	230	58	18	79	250
Aroclor-1260	44	430 J	LJO	30			
Total Metals (mg/Kg)				27000	27500	6430 J	14800
Aluminum	24400	28400	12600 J	26000	27500 117 J	1.7 J	4.2
Antimony	13.9 J	10.2 J	1.5 UJ	68.0 J	4550	15.9	186
Arsenic	875	670	121	3230	4550 52.5	63.8	107
Barium	53.4	126	149	78.8	1.7	0.45 J	1.1
Beryllium	1.7	2.1	0.97 J	2.0	11.4	3.1	3.8
Cadmium	12.4	19.9	9.5	11.7 8920	7160	3020	6950
Calcium	8590	9970	5800		726	61.2 J	562
Chromium	934	3580	303 J	2120 36.2	38.1	7.2 J	9.7
Cobalt	16.5	46.1	25.3 J	36.2 2080	1980	113	450
Copper	1870	2040	402	2080 90200	1980	12800 J	20200
Iron	31900	57800	31700 J		1220	340 J	771
Lead	559	787	606 J	1240	1480	1930	2370
Magnesium	1500	2410	2980	1860	1460 586	123	330
Manganese	409	760	728	776	44.8	0.25 J	14.3
Mercury	24.2	15.4	1.9	27.5	44.8 20.5	15.9 J	19.4
Nickel	18.4	50.1	32.8 J	28.3	20.3 450 U	715 J	741
Potassium	345 U	877	956 J	795	430 0	112 1	

TABLE 3-1. ANALYTES DETECTED IN 1997 SEDIMENT SAMPLES  $^{(1,2)}$ 

M&E SAMPLE ID:	SED-10-02	SED-10-04	SED-11-01	SED-12-01	SED-12-03	SED-13-01	SED-13-03
DAS SAMPLE NUMBER:	DAM352	DAM355	DAM336	DAM357	DAM358	DAM341	DAM342
DATE SAMPLED:	11/19/97	11/19/97	11/14/97	11/20/97	11/20/97	11/17/97	11/17/97
REMARKS:		FD of SED-10-01					
Selenium	11.5	8.3 J	2.5 J	18.3	30.3	0.53 J	3.1 J
Silver	0.57 J	0.17 J	0.92 J	0.54 J	1.2 J	0.21 J	0.32 J
Sodium	642 U	815	421	708 U	493 U	622	798
Thallium	1.3 U	2.6 U	0.44 UJ	1.2 U	1.5 U	0.43 UJ	0.43 UJ
⁄anadium	47.8	78.4	58.8	107	127	27.4	54.0
Zinc	1570	4700	2260 J	2640	4760	. 445 J	641 J
AVS-SEM (umoles/g)		•					0.50.1
Arsenic	4.53 J	0.80 J	2.98 J	3.40 J	2.99 J	0.05 U	0.50 J
Cadmium	0.05 U	0.12	0.10 UJ	0.05 U	0.08	0.05 U	0. <b>05</b> U
Copper	0.61	0.05 U	1.45 J	1.27	0.62	0.05 UJ	0. <b>59</b> J
Lead	1.87 J	1.53	1.29 J	1.54	3.26	1.17 J	2.79 J
Víckel	0.23 J	0.83 J	0.44 J	0.26 J	0.28 J	0.15 J	0.21 J
Sulfide	5.15	64.8	15.7	35.8	11.6	4.80	3.79
Zinc	9,56 J	25.0 J	17.8 J	13.0 J	13.3 J	5.79 J	4.96 J
ZINC SEM/AVS Ratio	2.38	0.42	1.34	0.45	1.51	1.48	2.26
TCO (mg/Kg)			•				*******
Total Combustible Organics	463000 J	459000 J	387000 J	912000 J	431000 J	232000 J	118000 J

reported in Appendix B.

- J Quantitation is approximate due to limitations identified in the quality control review
- U Value reported is the sample-specific detection limit
- R Value is rejected
- UJ Sample-specific detection limit is approximate due to limitations identified in the quality control review

<sup>(1)</sup> Validated data for all sediment and field QC samples collected during M&E's field investigation are

<sup>(2)</sup> Data presented for an analyte if the analyte was detected in at least one sediment sample

TABLE 3-1. ANALYTES DETECTED IN 1997 SEDIMENT SAMPLES (1,2)

M&E SAMPLE ID:	SED-15-01	SED-18-02	SED-18-03	SED-19-01	SED-20-01	SED-21-01	SED-21-05 DAM337
DAS SAMPLE NUMBER:	DAM340	DAM347	DAM348	DAM354	DAM339	DAM335	11/14/97
DATE SAMPLED:	11/17/97	11/18/97	11/18/97	11/19/97	11/17/97	11/14/97	FD of SED-21-01
REMARKS:					<u></u>	FD of SED-21-05	PD ()1 3ED-21-01
/OCs (ug/Kg)				22 111	46 UJ	19 J	22 J
Benzene	25 UJ	24 UJ	31 UJ	32 UJ	46 UJ 114 UJ	75 UJ	76 UJ
Chloroform	63 UJ	59 UJ	76 UJ	79 UJ	562 J	75 UJ	76 UJ
is-1,2-Dichloroethene	187 J	37 J	76 UJ	82 J	46 UJ	30 UJ	31 UJ
Ethylbenzene	25 UJ	24 UJ	31 UJ	32 UJ	· ·	75 UJ	76 UJ
m/p-Xylene	63 UJ	59 UJ	76 UJ	79 UJ	114 UJ	255 J	37 <b>9</b>
Naphthalene	1 <del>9</del> 0 UJ	178 U	229 U	237 UJ	343 U	233 J 225 UJ	229 U
Tetrachloroethene	190 UJ	178 U	229 U	237 UJ	790 U		229 U
trans-1,2-Dichloroethene	190 UJ	178 U	229 U	237 UJ	343 U	19 UJ 75 UJ	76 UJ
Trichloroethene	206 J	59 UJ	76 UJ	42 J	2025 J		70 UJ 229 U
Vinyl chloride	190 UJ	178 U	229 U	237 UJ	343 U	225 UJ	229 0
PAHs (ug/Kg)			•	n	11 J	140	
2-Methylnaphthalene	67 U	15 J	21 J	R	R	67 J	
Acenaphthene	R	30 J	29 J	R R	67 UJ	38 J	
Acenaphthylene	67 U	16 J	23 J		26 J	420	
Anthracene	50 J	150	130	18 J	210	390	
Benzo(a)anthracene	200	1400	1400	130 J	290 J	890 J	
Benzo(a)pyrene	180 J	1900	1900 J	160 J	330	980	
Benzo(b)fluoranthene	440 J	2500 J	2400 J	290 J	210 J	710 J	
Benzo(g,h,i)perylene	150 J	1400	1600 J	120 J	370	67 U	
Benzo(k)fluoranthene	270 J	1200 J	920 J	200 J	400	940	
Chrysene	410	2200	2000	230 J	400 97 J	240 J	
Dibenz(a,h)anthracene	60 J	430	410 J	46 J	720	1400	
Fluoranthene	990	4000	3100	410 J	720 29 J	240	
Fluorene	50 J	110	94	14 J	-	620 J	
Indeno(1,2,3-cd)pyrene	170 J	1400	1500 J	140 J	210 J	1300 J	
Naphthalene	15 J	29 J	34 J	12 J	14 J	600	
Phenanthrene	400	1400	1200	160 J	240		
Pyrene	590	3100	2600	340 J	540 J	1300	
PAH-SIMS (ug/Kg)					14 UJ		
Naphthalene	15 UJ			11 U 6 U	14 UJ 13 J		
2-Methylnaphthalene	10 J			6 U 1 J	6 J		
Acenaphthylene	7 J			1 J 3 J	6 J		
Acenaphthene	8 J 69 J			14 J	36 J		
Fluorene	б <del>у</del> Ј 34 Ј			18 J	45 J		
Dibenz(a,h)anthracene	34 J						

TABLE 3-1. ANALYTES DETECTED IN 1997 SEDIMENT SAMPLES (1,2)

M&E SAMPLE ID:	SED-15-01	SED-18-02	SED-18-03	SED-19-01	SED-20-01	SED-21-01	SED-21-05
DAS SAMPLE NUMBER:	DAM340	DAM347	DAM348	DAM354	DAM339	DAM335	DAM337
DATE SAMPLED:	11/17/97	11/18/97	11/18/97	11/19/97	11/17/97	11/14/9 <b>7</b>	11/14/97
REMARKS:						FD of SED-21-05	FD of SED-21-01
'esticides (ug/Kg)							
i,4'-DDD	6.9 J	23 J	4,9 J	9.0	8.4 J	35 J	
1,4'-DDE	3.8 J	28	30	3.2	4.4	22	
1,4'-DDT	1.8 J	1.2 J	1.7 J	0.33 U	4.6 J	4.4 J	
Aldrin	0.17 UJ	0.16 U	0.17 U	0.16 U	R	1.6 U	
alpha-BHC	0.17 UJ	0.16 U	0.17 U	0.16 U	0.16 U	1.6 U	
alpha-DhC alpha-Chlordane	6.0 J	7.7 J	17 J	0.98 J	12 J	29	
iepna-Cmoruane leita-BHC	0.17 UJ	0.16 U	0.17 U	0.16 U	0.16 U	1.8 J	
	0.57 J	0.33 J	0.94 J	2.5 J	0.70 J	3.2 U	
Dieldrin	0.45 J	0.16 U	0.17 U	0.44 J	0.16 U	1.6 U	
Endosulfan I	0.43 J 0.33 UJ	0.10 U	0.34 U	0.33 U	0.31 U	5.2	
Endosulfan II		0.33 U	0.41 J	0.33 U	0.31 U	3.2 U	
Endosulfan Sulfate	0.33 UJ	0.33 U 1.3 J	1.1 J	0.92 J	0.97 J	3.6 J	
Endrin	0.48 J	1.3 J 2.4 J	3.6 J	0.33 U	1.3 Ј	9.0 J	
Endrin Aldehyde	2.0 J			0.16 U	0.16 U	1.6 U	
gamma-BHC	0.17 UJ	0.16 U	0.17 U	0.16 U 0.59 J	6.8 J	19 J	
gamma-Chlordane	0.98 J	1.9 J	26 J			1.6 U	
Heptachlor Epoxide	0.17 J	0.16 U	0.17 U	0.16 U	0.49 J	1.0 0	
PCBs (ug/Kg)					20.1	240 1	
Aroclor-1248	43 J	60 J	170 J	35 J	28 J	240 J	
Aroclor-1260	83	90	190	11	79	420	
Total Metals (mg/Kg)							
Aluminum	6100 J	20300	24100	10100	13500 J	11300 J	
Antimony	3.9 J	8.9 J	9.2 J	5.2 J	3.8 J	2.0 J	
Arsenic	184	1180	839	182	236	258	
Barium	65.6	143	169	66.5	44.2	181	
Beryllium	0.76 J	1.8	2.3	2.9	1.2 J	0.93 J	
Cadmium	17.7	15.1	16.0	7.4	15.3	14.1	
Calcium	13400	- 7300	6860	11800	11100	8290	
Chromium	537 J	3890	4340	226	1090 J	5590 J	
Cobalt	23.1 J	44.3	54.9	18.7	28.3 J	26.9 J	
Copper	306	1010	1310	557	1060	613	
Iron	16000 J	99600	85100	19800	18500 J	20800 J	
Lead	157 J	469	678	197	450 J	546 J	
Magnesium	1300	2780	2930	1300	1530	2170	
Manganese	956	1630	1580	677	711	268	
Mercury	1.3 J	7.5	5.5	0.98	1.3 J	0.15	
Nickel	18.9 J	42.5	46.7	14.6	27.5 J	29.8 J	
Potassium	243 J	894	1050	342 U	454 J	531 J	

TABLE 3-1. ANALYTES DETECTED IN 1997 SEDIMENT SAMPLES (1,2)

M&E SAMPLE ID:	SED-15-01	SED-18-02	SED-18-03	SED-19-01	SED-20-01	SED-21-01	SED-21-05
DAS SAMPLE NUMBER:	DAM340	DAM347	DAM348	DAM354	DAM339	DAM335	DAM337
DATE SAMPLED:	11/17/97	11/18/97	11/18/97	11/19/97	11/17/97	11/14/97	11/14/97
REMARKS:	****					FD of SED-21-05	FD of SED-21-01
Selenium	1.2 UJ	8.0 J	9.0 J	5.6	3.3 J	3.2 J	
ilver	0.29 J	0.74 J	0.62 J	0.12 UJ	0.29 J	0.30 J	
odium	642	648 U	551 U	706 U	612	796	
Thallium	2.4 J	1.0 U	1.9 U	1.5 U	0.44 UJ	0.44 UJ	
/anadium	22.4	76.0	95.9	25.4	38.2	46.2	
Zinc	2490 J	4020	3600	1080	2380 J	4940 J	
VS-SEM (umoles/g)							
Arsenic	0.47 J	3.43 J	3.00 J	0.52 J	1.12 J	0.19 J	
Cadmium	0.20 J	0.11	0.08	0.05 U	0.09 J	0.09 UJ	
Соррег	0.05 UJ	2.18	4.11	0.38	1.29 J	0.05 UJ	
Lead	0.81 J	1.70	2.38	0.60	1.43 J	1.68 J	
Vickel	0.52 J	0.47 J	0.61 J	0.15 J	0.72 J	0.17 J	
Sulfide	31.3	17.1	15.9	6.88	17.7	28.0	
Zinc	26.1 J	19.5 J	18.9 J	9.55 J	20.6 J	25.7 J	
SEM/AVS Ratio	0.88	1.40	1.64	1.55	1.36	0.99	
TCO (mg/Kg)						•	
Total Combustible Organics	702000 J	361000 J	423000 J	686000 J	441000 J	344000 J	

(1) Validated data for all sediment and field QC samples collected during M&E's field investigation are reported in Appendix B.

(2) Data presented for an analyte if the analyte was detected in at least one sediment sample

- J Quantitation is approximate due to limitations identified in the quality control review
- U Value reported is the sample-specific detection limit
- R Value is rejected
- UJ Sample-specific detection limit is approximate due to limitations identified in the quality control review

TABLE 3-1. ANALYTES DETECTED IN 1997 SEDIMENT SAMPLES (1,2)

	SED-22-02	SED-24-03	SED-25-02
M&E SAMPLE ID:	DAM321	DAM320	DAM346
DAS SAMPLE NUMBER:	11/12/97	11/12/97	11/18/97
DATE SAMPLED:	11/12/9/		
REMARKS:			
VOCs (ug/Kg)		9 ()]	4 J
Benzene	27 UJ	23 UJ	12 UJ
Chloroform	68 UJ	23 UJ	12 UJ
cis-1,2-Dichloroethene	73 J	23 UJ 9 UJ	5 UJ
Ethylbenzene	27 UJ	23 UJ	. 12 UJ
m/p-Xylene	68 UJ	23 UJ 69 U	36 UJ
Naphthalene	205 U	69 U	36 UJ
Tetrachloroethene	3164		36 UJ
trans-1,2-Dichloroethene	205 U	69 U	12 UJ
Trichloroethene	803	23 UJ 69 U	36 UJ
Vinyl chloride	205 U	69 0	30 03
PAHs (ug/Kg)			22 1
2-Methylnaphthalene	11 J	23 J	22 J 69 J
Acenaphthene	R	50 J	*
Acenaphthylene	8 J	44 J	22 J
Anthracene	46 J	250	330
Benzo(a)anthracene	300	960	1300
Benzo(a)pyrene	200	790 J	1300
Benzo(b)fluoranthene	440	880	1500 J
Benzo(g,h,i)perylene	190	550 J	650
Benzo(k)fluoranthene	290	400	740 J
Chrysene	400	920	1400
Dibenz(a,h)anthracene	44 J	120 J	220
Fluoranthene	840	2600	2500
Fluorene	23 J	130	120
Indeno(1,2,3-cd)pyrene	220	620 J	670
Naphthalene	24 J	23 J	32 J
Phenanthrene	490	1300	1600
Pyrene	610	2400	2300
PAH-SIMS (ug/Kg)			
PAH-SIMS (ug/Kg) Naphthalene	25 J		
Naphtnaiene 2-Methylnaphthalen <del>e</del>	13 J		
Acenaphthylene	12 J		
Acenaphthene	11 J		
Fluorene	26 J		
Dibenz(a,h)anthracene	36 J		

TABLE 3-1. ANALYTE'S DETECTED IN 1997 SEDIMENT SAMPLES (1,2)

			SED-25-02
M&E SAMPLE ID:	SED-22-02	SED-24-03	DAM346
DAS SAMPLE NUMBER:	DAM321	DAM320	
DAS SAMPLE HOMBER:	11/12/97	11/12/97	11/18/97
DATE SAMI LED. REMARKS:			
esticides (ug/Kg)		390	11 J
.4'-DDD	9.6 J	***	4.4 J
,4'-DDE	31	37	2.1 J
,4'-DDT	2.7 ]	24 J	0.16 U
,4°-001   drin	0.15 U	1.6 U	0.16 U
viariu slpha-BHC	0.15 U	1.6 U	1.3 J
alpha-Bric Alpha-Chlordane	1.2 J	2.0 J	0.16 U
	0.16 J	1.6 U	0.32 U
delta-BHC	0.38 J	3.2 U	0.32 U
Dieldrin	0.18 J	1.6 U	•
Endosulfan I	0.48 J	3.2 U	0.33 J
Endosulfan II	0.31 U	3.2 U	0.32 U
Endosulfan Sulfate	0.31 U	3.2 U	0.98 J
Endrin	3.3 J	5.9 J	4.4 J
Endrin Aldehyde	0.15 U	1.6 U	0.16 U
gamma-BHC	0.19 J	4.2	1.3 J
gamma-Chlordane	0.19 J	1.6 U	0.36 J
Heptachlor Epoxide	0.27		
PCBs (ug/Kg)	8.1 J	290 J	56 J
Arocior-1248	29	200	47
Aroclor-1260			
Total Metals (mg/Kg)	2700 J	8440 J	7190
Aluminum	8.0 J	1.2 J	0.64 J
Antimony	6.6	30.7	5.4
Arsenic	32.9	43.6	48.5
Barium	0.48 J	0.80 1	0.37
Beryllium	0.61	2.0	0.78 J
Cadmium	19600	2410	2150
Calcium	11.7 J	410 J	26.1
Chromium	2.7 J	12.1 J	8.1
Cobalt	26.0	130	52.7
Copper	2710 J	13200 J	14700
Iron	3970 J	566 J	210
Lead	1370	2880	3490
Magnesium	36.7	131	190
Manganese	0.50	0.71	0.10 14.6
Mercury	9,2 J	23.7 J	14.6 1210
Nickel	260 J	1170 J	1210
Potassium			32

TABLE 3-1. ANALYTES DETECTED IN 1997 SEDIMENT SAMPLES (1,2)

SED-22-02	SED-24-03	SED-25-02
8EB 22 0=	DAM320	DAM346
	11/12/97	11/18/97
11/12/97		
8.1 J	•	0.48 UJ
0.22 J		0.20 J
546		187 U
0.44 UJ	0.12 UJ	0.14 U
38.8	80.3	29.7 J
34.2 J	251 J	178
	0.05.11	0.05 U
0.05 U	0.00	0.05 U
0.05 U	0.05	0.4-
0.53 J	0.05 UJ	0.11
20.4 J	1. <b>7</b> 9 J	0.90
0.35 J	0.38 J	0.05 J
0.050 U	10.2	2.03
2.10 J	3.69 J	1.84 J
467.6	0.57	1.43
602000 I	153000 I	59200 J
	0.22 J 546 0.44 UJ 38.8 34.2 J 0.05 U 0.05 U 0.53 J 20.4 J 0.35 J 0.050 U 2.10 J	DAM321 DAM320 11/12/97 11/12/97  8.1 J 0.91 J 0.22 J 2.5 J 546 516 0.44 UJ 0.12 UJ 38.8 80.3 34.2 J 251 J  0.05 U 0.05 U 0.05 U 0.05 U 0.53 J 0.05 UJ 1.79 J 0.35 J 0.38 J 0.050 U 10.2 2.10 J 3.69 J 467.6 0.57

(1) Validated data for all sediment and field QC samples collected during M&E's field investigation are reported in Appendix B.

(2) Data presented for an analyte if the analyte was detected in at least one sediment sample

- J Quantitation is approximate due to limitations identified in the quality control review
- U Value reported is the sample-specific detection limit
- R Value is rejected
- UJ Sample-specific detection limit is approximate due to limitations identified in the quality control review

# 4.1 Comparison of 1995 and 1997 Sediment Data

The 1995 and 1997 sediment data were compared with each other in order to identify any differences between data sets in terms of the analytical methods employed, the locations where samples were collected, and the time frame between collection of samples. This is helpful in understanding how the data can be used together and if any time or spatial trends are important.

Tables 4-1 and 4-2 summarize the 1995 and 1997 sediment data in terms of the analytes detected, the range of detected values, the sample site and locations where the maximum detected value was found, the frequency of detection, and, where applicable, the range of sample-specific detection limits. The data sets used for both years contained only the locations that were sampled by M&E in 1997. Both M&E's and FW's data were handled in the same manner. The treatment of data used to obtain the summary tables is described in Appendix E. In comparing the tables, it should be noted that for some parameters, nondetected data were rejected for certain 1995 sample locations. This can be seen in the frequency of detection, when fewer than 28 sample locations are reported for the total number of locations.

Similar to the 1995 data, PAHs were the most prevalently detected organic compounds in 1997 and were detected in higher concentrations than other organic compounds. Although PAHs were detected at all of the 1997 sample locations compared to most of the 1995 sample locations, the range of concentrations detected between the two data sets is similar. The highest individual concentrations were detected for more PAHs at the SD-07 site in both data sets. Also, the highest detected concentrations tended to occur with the higher molecular weight PAHs in both data sets. In addition to PAHs, a few other semivolatile organic compounds (SVOCs) were detected in 1995. Due to their prevalence at the site and their associated risks, PAHs were the only SVOCs analyzed for in 1997.

Pesticides were also frequently detected in both the 1995 and 1997 data sets. Like PAHs, pesticides were detected at more sample locations in 1997 compared to 1995. The frequency of detection was above 70% for eight pesticides in the 1997 data set. Many of the same pesticides were detected in both data sets, with the DDT and chlordane groups being the predominant types in terms of frequency of detection and concentrations.

PCB Aroclor 1260 was detected in two locations during FW's 1995 sampling effort. In 1997, Aroclors 1248 and 1260 were detected in all 28 of the locations sampled, including the two reference locations. Comparing the two data sets, the range of concentrations detected are not substantially different between each set. However, the widespread detection of Aroclor 1260 may be attributed to the lower detection limit for the DAS analyses in 1997 relative to the RAS analyses conducted in 1995.

Although relatively few volatile organic compounds (VOCs) were also detected during both sampling periods, the types of VOCs reported differed. For instance, in 1995 ketones were found in relatively high concentrations compared to the three chlorinated VOCs that were detected.

TABLE 4-1. DATA SÚMN	MARY FOR A	ANALYTES DE		Eranjana	Range of	SSDLs
I MADESTS & 13 25	Range of Det	C.C. C. C. C.		of Detection	Minimum	Maximum
Parameter/Analyte	Minimum	Maximum	Location	OF Become	:::::::::::::::::::::::::::::::::::::::	
	::::::::::::::::::::::::::::::::::::::	,		'		
Volatile Organic Compounds (	<u>ug/Kg)</u>	5.00	SD-20-01-ME	18 / 26	9	225
cis-1,2-Dichloroethene	7	562	SD-11-01-ME	1 / 26	28	343
trans-1,2-Dichloroethene	387	387	SD-20-01-ME	14 / 26	9	114
Trichloroethene	6	2,025	SD-20-01-ME SD-22-02-ME	1 / 26	2.8	343
Tetrachloroethene	3,164	3,164	SD-22-02-ME SD-11-01-ME	1 / 26	28	343
Vinyl chloride	255	255	SD-11-01-ME SD-21-01-ME	8 / 26	4	225
	6	22		4 / 26	4	46
Benzene	5	9	SD-06-03-ME	2 / 26	9	114
Ethylbenzene Xylene, m/p-	10	25	SD-06-03-ME	2,20		
				1	1	
Polycyclic Aromatic Hydroca	bons (ug/Kg)	(4)	on 21 01 ME	26 / 26	1	67
Naphthalene	12		SD-21-01-ME	24 / 26	1	67
Naphthalene 2-Methylnaphthalene	10	140	SD-21-01-ME	26 / 26	1	67
	3	230	SD-07-10-ME	1	1	67
Acenaphthene	1	170	SD-04-02-ME	<u> </u>	1	67
Acenaphthylene	14	: 640	SD-07-02-ME	1	67	67
Fluorene	160	12,000	SD-07-10-ME	1	67	67
Phenanthrene	18	890	SD-07-10-ME		67	67
Anthracene	410	23,000	SD-07-10-ME		67	67
Fluoranthene	340	15,000	SD-07-10-ME		67	67
Pyrene	130	9,600	SD-07-10-ME	26 / 26	67	67
Benzo(a)anthracene	230	10,000	SD-07-10-ME		67	67
Chrysene	290	16,000	SD-07-10-MI		67	67
Benzo(b)fluoranthene	200	4,100	SD-07-02-MI			67
Benzo(k)fluoranthene	l.	10,000	SD-07-10-MI	36 / 26	67	67
Benzo(a)pyrene	160	2,000	SD-07-10-M	E 26 / 26	2	67
Dibenz(a,h)anthracene	44	5,300	SD-07-10-M	E 26 / 26	67	67
Benzo(g,h,i)perylene	120	( 000	SD-07-10-M		67	01
Indeno(1,2,3-cd)pyrene	140	5,900	1			1
- constitution		:		E 26 / 26	0.31	3.9
Pesticides (ug/Kg)	1.9	310	SD-13-03-M		0.31	
4,4'-DDD	0.9	160	SD-10-01-M		0.31	2.0
4,4'-DDE	1.1	21	SD-04-02-M		0.15	
4,4'-DDT	0.17	0.17	SD-01-07-M	1.12		
alpha-BHC	0.16	25	SD-06-03-N			
delta-BHC	0.23	0.34	SD-12-03-N	·	1	
gamma-BHC (Lindane)	0.98	0.2	SD-13-03-N	ИЕ 26 / 26 ИЕ 26 / 26	1	
alpha-Chlordane	0.19	7 CA	SD-13-03-1		1 .	• :
gamma-Chlordane	0.21	• • •	SD-10-01-I	ME 3 / 25		
Aldrin	0.33		SD-10-01-	ME 19 / 26	· 1	*
Dieldrin	0.48	1.77	SD-06-03-1	ME 22 / 26		
Endrin	0.8	,	SD-07-02-	ME 20 / 20	· .	
Endrin Aldehyde	0.1		5 SD-15-01-	ME 4 / 20	-	_
Endosulfan I	L	·	SD-21-01-	ME 9 / 2		-
Endosulfan II	0.2	•	8 SD-01-07-	ME 7 / 2		
Endosulfan Sulfate	0.3		1 40 01	ME 8 / 2	6 0.1	1.5
Heptachlor Epoxide	0.1	,				
		*		201	,	.6 3
PCBs (ug/Kg)	8.	1 56	O SD-10-01	-ME 26 / 2		.6 3
Aroclor 1248			O SD-21-01			

TABLE 4-1. DATA SUMMARY FOR ANALYTES DETECTED IN THE 1997 SEDIMENT SAMPLES (1.2.3)

TABLE 4-1. DATA SUMI	MARY FUR A	MATERIANA Mariana Maria Cana	Maximum	Frequency		
	Range of Dete		Location	of Detection	Minimum	Maximum
arameter/Amalyte	Minimum	Maximum		<b></b>	}::=::::::::::::::::::::::::::::::::::	
			1			
			1			
<u> Jetals - Low Concentration (r</u>	ng/Kg)	20.250	SD-10-01-ME	26 / 26	3.5	8.2
Aluminum	2,700	29,250	SD-12-03-ME	25 / 26	0.067	0.46
Antimony	1.4	117	SD-12-03-ME	26 / 26	0.22	0.51
Arsenic	6.6	4,550	SD-21-01-ME	26 / 26	0.018	0.043
Barium	33	181	h	26 / 26	0.027	0.064
Beryllium	0.295	2.9	SD-19-01-ME	26 / 26	0.0048	0.81
Cadmium	0.61	19	SD-04-02-ME	26 / 26	1.6	3.8
Calcium	1,340	19,600	SD-22-02-ME	26 / 26	0.055	0.13
Chromium	12	5,590	SD-21-01-ME	26 / 26	0.37	0.88
Cobalt	2.7	55	SD-18-03-ME	26 / 26	0.082	0.19
Copper	26	2,315	SD-10-01-ME		0.57	1.3
Iron	2,710	107,000	SD-12-03-ME	26 / 26	0.26	0.62
-	157	3,970	SD-22-02-ME	26 / 26	0.82	8.1
Lead	1,300	5,330	SD-07-05-ME	26 / 26	0.018	0.043
Magnesium	37	1,630	SD-06-03-ME	26 / 26	1	0.0059
Manganese	0.15	45	SD-12-03-ME	26 / 26	0.0037	0.34
Mercury	9.2	47	SD-18-03-ME	26 / 26	0.15	6.3
Nickel	243	1.620	SD-07-05-ME	23 / 26	2.7	0.69
Potassium	0.53	30	SD-12-03-ME	24 / 26	0.052	
Selenium	0.33	1.2	SD-12-03-ME	25 / 26	0.0094	0.043
Silver	94	798	SD-13-03-ME	16 / 26	0.22	0.51
Sodium	1	2.4	SD-15-01-ME	1 / 26	0.038	0.17
Thallium	2.4	127	SD-12-03-ME	26 / 26	0.073	0.17
Vanadium	22	5,170	SD-02-02-ME	26 / 26	0.16	0.39
Zinc	34	5,170	0.0	İ		
l .	1		AWS/SEM (mo/l	(p)		
Acid Volatile Sulfide/Simult	taneously Extra	cted ivietais	SD-02-02-ME	25 / 26		
Sulfide	124	7,172	SD-10-02-ME	4. (0/		
Arsenic	3.7	339	SD-15-01-ME	1		
Cadmium	5.6	22	SD-13-01-ME	1		
Copper	4.0	261	SD-22-02-ME	'   "		
Lead	124	4,227	SD-03-02-ME			
Nickel	4.7	58	1	1		
Zinc	138	2,419	SD-02-02-MI	20 1 20		
SEM/AVS Ratio (umoles/	/g) 0.22	468	SD-22-02-MI	26 / 26		
Total Combustible Organic Total Combustible Organ	  cs (mg/K.g)  ics  59,100	912.00	0 SD-12-01-M	E 26 / 26	2,00	0 ; 2,00

- 1. M&E's analytical data are presented in Appendix B and detected data are summarized in Table 3-1. Field NOTES: duplicate samples were averaged together as described in Appendix E prior to incorporation into summaries.
- 2. Data summaries were developed assuming positive detections and included approximated values (due to validation qualifiers) and included approximated values less than SSDLs.
- 3. The two reference locations that were re-sampled are not included in the data set (SD-24 and SD-25).
- 4. PAH results combined from both full scan and SIM analyses.

SSDLs - Sample-Specific Detection Limits

<sup>--</sup> Not applicable

TABLE 4-2. DATA SUMMARY FOR ANALYTES DETECTED IN THE 1995 SEDIMENT SAMPLES (1,2,3)

			1 1	P	Ranj	d Conc. (4)
	Range of Det		Maximum	Frequency	Minimum	Maximum
arameter/Analyte	Minimum	Maximum	Location	of Detection (3)	Minimum	
olatile Organic Compounds	(ug/Kg)				12	29
1,2-Dichloroethene(total)	9	29	SD-15-01-FW	4 / 17	12	29
Tetrachloroethene	120	120	SD-22-02-FW	1 / 14		56
Methylene Chloride	28	28	SD-22-02-FW	1 / 14	12	26
2-Butanone	9	290	SD-19-01-FW	11 / 19	12	430
Acetone	210	590	SD-20-01-FW	4 / 18	16	430
1 1 11 Chamania Campro	unde (uø/Kø)					
emivolatile Organic Compo	83	160	SD-04-03-FW	3 / 14	410	2,900
Naphthalene	130	130	SD-07-10-FW	1 / 14	410	2,900
2-Methylnaphthalene	250	480	SD-07-10-FW	3 / 14	410	2,900
Acenaphthylene	i	340	SD-04-03-FW	6 / 15	410	2,900
Acenaphthene	100	530	SD-07-05-FW	7 / 15	410	890
Fluorene	180	6,300	SD-06-03-FW	14 / 18	410	890
Phenanthrene	74	•	SD-06-03-FW	10 / 15	410	890
Anthracene	75	1,300	SD-07-05-FW	21 / 23	410	890
Fluoranthene	210	15,000	SD-07-05-FW	20 / 22	410	870
Pyrene	100	12,000		16 / 19	410	890
Benzo(a)anthracene	100	6,400	SD-07-05-FW	1	410	890
Chrysene	150	9,800	SD-07-05-FW		410	870
Benzo(b)fluoranthene	120	12,000	SD-07-05-FW		410	870
Benzo(k)fluoranthene	110	14,000	SD-07-05-FW	18 / 21	l.	890
Benzo(a)pyrene	130	6,300	SD-07-05-FW		410	2,900
Dibenz(a,h)anthracene	180	1,100	SD-04-03-FW		410	890
Benzo(g,h,i)perylene	140	2,300	SD-04-03-FW	I .	410	
Indeno(1,2,3-cd)pyrene	79	2,600	SD-07-05-FW		410	890
Dibenzofuran	87	170	SD-04-03-FW		410	2,900
Bis(2-ethylhexyl)phthalate	140	13,000	SD-07-05-FW		410	3,300
Butylbenzylphthalate	230	620	SD-07-05-FW	3 / 16	410	2,900
Di-n-octylphthalate	430	430	SD-07-05-FW	1 / 15	410	2,900
	100	560	SD-10-01-FW		410	2,900
N-Nitrosodiphenylamine Carbazole	130	580	SD-07-05-FW	7 / 16	410	2,900
Pesticides (ug/Kg)	1 22	140	SD-04-03-FW	/ 16 / 22	4.1.	43
4,4'-DDD	2.3	125	SD-10-01-FV		4.1	43
4,4'-DDE	1.7	47	SD-07-10-FV		4.1	49
4,4'-DDT	0.45	1	SD-07-02-FV	·	2.1	25
alpha-BHC	0.32	2.7	SD-07-02-FV		2.1	25
beta-BHC	1.5	3.9	I I	9	2.1	25
delta-BHC	1.8	4.2	SD-07-02-FV	1	2.1	25
gamma-BHC (Lindane)	1.4	1.4	SD-07-02-FV	L.	2.1	22
alpha-Chlordane	1.6	60	SD-21-01-FV		2.1	25
gamma-Chlordane	0.34	36	SD-07-05-FV			25
Aldrin	0.47	1.6	SD-13-03-FV	1	2.1	49
Dieldrin	1.3	5.7	SD-07-05-FV		4.1	49
Endrin	0.2	. 16	SD-21-01-FV		4.1	
Endrin aldehyde	1	10	SD-13-03-FV		4.1	49
Endrin ketone	2.3	2.3	SD-07-02-FV		4.1	49
Endosulfan I	0.62	68	SD-21-01-F		2.1	2.5
Endosulfan II	0.69	5.2	SD-07-02-F	w 3 / 14	4.1	49
	2.1	2.1	SD-07-05-F		4.1	49
Endosulfan sulfate	0.18	0.29	SD-07-05-F	L. L.	2.1	25
Heptachlor	i	3.7	SD-07-05-F		2.1	25
Heptachlor epoxide	1	12	SD-07-10-F	l.	21	250
Methoxychlor	6.3	1.4	U-10-1		1	

TABLE 4-2. DATA SUMMARY FOR ANALYTES DETECTED IN THE 1995 SEDIMENT SAMPLES (1,2,3)

TABLE 4-2. DATA SUN			,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,		Range of	
	Range of Det	ected Conc.	Maximum	Frequency		d Conc. (4)
arameter/Analyte	Miningum	Maximum	Location	of Detection (3)	Minimum	Maximum
	624					
CBs (ug/Kg)			an 10 02 FW	2 / 13	41	490
Aroclor 1260	17	310	SD-13-03-FW	2 / 13	7.	4,70
norganics (mg/Kg)						
Aluminum	3,560	30,200	SD-13-01-FW	26 / 26		. 2.1
Antimony	0.33	61	SD-13-01-FW	20 / 23	0.42	2.1
Arsenic	3.6	4,210	SD-13-01-FW	25 / 26	3.7	3.7
Barium	19	155	SD-13-03-FW	26 / 26		0.22
Beryllium	0.3	1.7	SD-10-02-FW	20 / 24	0.11	0.32
Cadmium	0.2	38	SD-01-06-FW	24 / 26	0.035	5.7
Calcium	1,300	15,000	SD-15-01-FW	26 / 26		
Chromium	7.5	2,335	SD-19-01-FW	26 / 26		
Cobalt	1.4	45	SD-19-01-FW	26 / 26		••
Copper	10	2,340	SD-13-01-FW	26 / 26		
Iron	2,310	163,500	SD-19-01-FW	26 / 26		
Lead	5.7	1,270	SD-13-01-FW	26 / 26		
Magnesium	715	7.610	SD-13-03-FW	26 / 26		
-	34	1,400	SD-15-01-FW	26 / 26		
Manganese	0.063	20	SD-13-01-FW	23 / 26	0.021	0.1
Mercury	4.4	53	SD-13-03-FW	26 / 26		~~
Nickel	195	2.030	SD-13-03-FW	22 / 25	369	610
Potassium	2.3	26	SD-13-01-FW	12 / 24	0.43	2.7
Selenium	0.87	2	SD-13-01-FW	_	0.22	0.79
Silver	141	1,300	SD-18-03-FW		37	404
Sodium	0.47	3.8	SD-07-10-FW		1.2	1.6
Thallium	12	120	SD-13-03-FW	1		
Vanadium	23	7,380	SD-01-06-FW			
Zinc	2.5	7,500				
Cyanide	. 1.8	1.8	SD-07-05-FW	1 / 15	0.59	1.6
Acid Volatile Sulfide/Simult	 aneously Extr	acted Metals	AVS/SEM (mg/	ı <u>Kg</u> )		
Sulfide	13	13,482	SD-07-05-FW	26 / 26		
Cadmium	0.23	34	SD-15-01-FW	24 / 26	0.1	0.37
Copper	42	2,270	SD-13-01-FW	20 / 26	12.3	138
	7.4	1,590	SD-13-01-FW	25 / 26	34	34
Lead	0.18	1	SD-13-03-FW	2 / 26	0.003	0.02
Mercury Nickel	39	258	SD-19-01-FW	*	2.3	221
INICKCI	"					
SEM/AVS Ratio (umoles/g	<u>;</u> 0	70	SD-13-03-FW	26 / 26	**	
Total Combustible Organic	l s/Total Organ	ic Carbon (mg	/Kg)			:
Total Combustible Organic		432,000	SD-15-01-FV			
Total Organic Carbon	6,525	288,000	SD-21-01-FV	v 7/7		

- 1. FW's analytical data are presented in the Preliminary Data Compendium (1996). Field duplicate samples were averaged together as described in Appendix E prior to incorporation into summaries. The sample group only includes sample station/locations that were re-sampled by M&E.
- 2. Data summaries were developed assuming positive detections and include approximated values.
- 3. The two reference locations that were re-sampled are not included in the data set (SD-24 and SD-25).
- 4. Sample-specific detection limits were not available for the data; therefore, nondetected values are shown.
- -- Not applicable or the nondetected values are not available because the analyte was detected in all samples in the data set.

In contrast, five chlorinated VOCs were reported in 1997 and concentrations were higher than those found in 1995. In addition, three aromatic VOCs were found in 1997 (benzene, ethylbenzene, and xylene) but were not reported in any 1995 sample locations.


Heavy metals were detected in over 75% of the sample locations in both 1995 and 1997 including antimony, arsenic, cadmium, chromium, cobalt, copper, lead, manganese, mercury, nickel, vanadium, and zinc. In general, concentration ranges for individual metals did not substantially differ between the two data sets. In contrast, the 1995 data exhibited higher AVS/SEM metal concentrations for copper and zinc compared to the 1997 data. Also, arsenic and zinc were reported by AVS/SEM at most of the sample locations in 1997, but not found at any locations in 1995.


Relatively elevated TCO levels were found in both the 1995 and 1997 data sets. The maximum TCO level reported in 1997 was greater than twice the level reported in 1995.


#### 4.2 Data Trends

To discuss data trends throughout the study area, data are discussed in terms of organic chemical groups and metals. The predominant types of organics detected consist of PAHs, phthalate compounds, pesticides, PCBs, and VOCs (aromatic, chlorinated, and ketones). For each organic group, detected concentrations for analytes with a related chemical type of that group are summed together to provide a total concentration for that organic group. Figure 4-1 presents total concentrations by sediment site/location for PAHs, pesticides, and PCBs, while Figure 4-2 presents total concentrations for aromatic VOCs, chlorinated VOCs, and ketones. The four heavy metals that were selected to represent trends include arsenic, chromium, lead, and mercury, which are shown on Figure 4-3.

Overall, PAHs and pesticides were detected in two or more sediment locations within each of the 28 sites sampled. PAHs were found in over 90% of the locations and pesticides were found in over 85% of the sediment locations sampled. Generally, sediment sites and locations exhibiting highest total PAH concentrations tended to have the more elevated total pesticide concentrations. The highest PAH and pesticide concentrations tended to occur at SED-07 compared to other sites. As reported by FW (1996), the physical characteristics of this site (a spillway, dam, and sand bar) contribute to being an area subject to accumulation of sediments and thus contaminants. Other sites with consistently elevated total PAH and pesticide concentrations include SED-04, SED-06, SED-13, and SED-18. Sites SED-04 and SED-06 are both described as depositional areas near the inlets to water bodies (i.e., Upper Mystic Lake). The predominance of these types of organics is expected given the close proximity of many of the locations to these types of features. The widespread use of pesticides (particularly the DDT and chlordane groups), along railroad lines and public roads as well as water bodies and public areas, was common practice prior to the 1980s. Run-off from these areas would be expected to transport both PAHs and pesticides into the water bodies. These organics would have the tendency to strongly bind to sediment and organic matter, and be further transported







downgradient, accumulating in depositional areas and bodies of water with little to no current.

As previously discussed, PCBs were found at all sediment locations sampled in 1997, but only two locations in 1995. The higher concentrations of PCBs detected in 1997 tended to occur at sediment locations exhibiting more elevated concentrations of PAHs and pesticides. The greater frequency of detections in 1997 may be attributed to the lower sample quantitation limits provided by the DAS Analytical Specifications.

In comparison to the above organic compounds, VOCs were less frequently detected during both sampling events, and when detected, concentrations were relatively low. In 1997, chlorinated VOCs were found at 10 of the 12 sites sampled. Sampling site SED-22 exhibited the highest total chlorinated VOC concentrations detected in both 1995 and 1997. Aromatic VOCs on the other hand were found at only 6 of the 12 sites sampled in 1997. Ketones (primarily acetone and 2-butanone) were reported in 11 of the 27 sites sampled in 1995. The majority of the VOCs detected during both sampling events were found in the vicinity of the Well G and Well H locations.

Metals exhibited a pattern of distribution similar to PAHs and pesticides. The range of concentrations between the two data sets is similar. More elevated concentrations of heavy metals tended to occur near roadways and rail lines and were characteristic of depositional areas. Sampling site SD-13 provided the greatest number of maximum metals concentrations in 1995 while the 1997 data did not show one location having a great deal more metals contamination than another location.

Overall, the sediment sampling at the Wells G&H Superfund Site verified the results of the previous sampling performed by FW in 1995. In addition, data gaps were filled using sampling techniques and DAS Analytical Methods specifically written for this site. The Data Quality Objectives and completeness goals were met by the sampling and analysis procedures provided in the QAPP. Sufficient data now exists to develop a site conceptual model and perform a baseline human health and ecological risk assessment for the Aberjona River System within the study area.

#### 5.0 REFERENCES

- Foster Wheeler Environmental Corporation (FW). 1996a. Preliminary Data Compendium -- Wells G&H RI/FS, Operable Unit 3, Aberjona River Study, Sampling Period July 31, 1995 to September 12, 1995. Prepared for the U. S. Environmental Protection Agency. 1996.
- Metcalf & Eddy (M&E). 1997a. Revised Draft Work Plan for Remedial Investigation/Feasibility Study (Risk Assessment) --- Aberjona River (Operable Unit 3), Wells G&H Superfund Site, Middlesex County, Massachusetts. Prepared for the U.S. Environmental Protection Agency. September 1997.
- Metcalf & Eddy (M&E). 1997b. Final Quality Assurance Project Plan for Wells G&H Superfund Site, Aberjona River Study (Operable Unit 3): Sediment Sampling for Human Health and Ecological Risk Assessments, Woburn, Massachusetts. Prepared for the U. S. Environmental Protection Agency. October 1997.
- U.S. Environmental Protection Agency (U.S. EPA). 1989a. Risk assessment guidance for Superfund. Volume II: Environmental evaluation manual. Part A. Interim Final. Office of Emergency and Remedial Response, Washington, D.C. EPA/540/I-89/001. March 1989.
- U.S. Environmental Protection Agency (U.S. EPA). 1989b. Risk assessment guidance for Superfund. Volume I: Human health evaluation manual. Part A. Interim Final. Office of Emergency and Remedial Response, Washington, D.C. EPA/540/I-89/002. December 1989.
- U.S. Environmental Protection Agency (U.S. EPA). 1989c. Supplemental risk assessment guidance for the Superfund program. Draft Final. Risk Assessment Work Group --- U.S. Environmental Protection Agency, Region I. EPA/901/5-89/001.
- U.S. Environmental Protection Agency (U.S. EPA). 1996. Region I, EPA-New England Data Validation Functional Guidelines for Evaluating Environmental Analyses. December 1996.